

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 242 855
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87105854.1

(51) Int. Cl.³: **A 61 K 7/48**
A 61 K 9/00, A 23 L 1/03

(22) Date of filing: 21.04.87

(30) Priority: 21.04.86 JP 91769/86

(43) Date of publication of application:
28.10.87 Bulletin 87/44

(84) Designated Contracting States:
AT BE CH DE ES FR GB IT LI NL SE

(71) Applicant: **LION CORPORATION**
3-7, Honjo 1-chome
Sumida-ku Tokyo(JP)

(72) Inventor: **Nagaoka, Yoshio**
No. 1-17-9-102, Higashiyotsugi Katsushika-ku
Tokyo(JP)

(72) Inventor: **Yasumasu, Tomoko**
No. 500-1-3-309, Hongo-cho
Funabashi-shi Chiba-ken(JP)

(72) Inventor: **Noda, Seiji**
No. 531-464, Wakamatsu-cho
Chiba-shi Chiba-ken(JP)


(72) Inventor: **Higo, Moriaki**
No. 6-8-8, Makuharinishi
Chiba-shi Chiba-ken(JP)

(74) Representative: **Kuhnen, Wacker & Partner**
Schneggstrasse 3-5 Postfach 1729
D-8050 Freising(DE)

(54) Crosslinking reaction inhibitor for collagen.

(57) The present invention relates to a crosslinking inhibitor which inhibits the formation of crosslinking (aged crosslinking) of collagen by utilizing the Maillard reaction between collagen and sugar. The inhibitor comprises a silicon compound having at least one silanol group in its molecule, such as methyl silanetriol, dimethyl silanediol, trimethyl silanol, dodecyl silanetriol, cyclohexyl silanetriol or propenyl silanetriol.

FIG. 1



SPECIFICATION

Title of the Invention

Crosslinking Reaction Inhibitor for Collagen

5

Background of the Invention

(1) Field of the Invention

10

The present invention relates to a crosslinking inhibitor which inhibits the formation of crosslinking (aged crosslinking) of collagen by utilizing the Maillard reaction between collagen and sugar.

(2) Prior Art

15

20

25

It has been previously known that amino acids added to foods and medicines, such as lysine, glutamic acid and alanine, react with sugars such as galactose and dextrose to produce a brown substance, this reaction being called the Maillard reaction. Techniques for inhibiting the above-described browning and inhibiting the production of discoloration over long-term preservation period have therefor been developed. For example, Japanese Patent Un-examined Publication No. 129945/1983 discloses a technique in which a thin film cover of a particular adhesive substance is provided on the periphery of at least one of granular sugar and amino acid and Japanese Patent Un-examined Publication No. 164683/1983 discloses a composition to which a polysaccharide containing 3% or less of water is added in an amount of 40% or more with respect to an amino acid.

On the other hand, evidence has been found that the hemoglobin of a persons afflicted with diabates mellitus and the collagen of aged persons are subjected to the Maillard reaction and the correlation between the Maillard reaction and diabetes or senescence has attracted considerable attention (with regard to diabetes, S. Rahbar; Chin. Chim. Acta, 22, 296 (1968) and with regard to senescence, D. Fujimoto; Biomedical Res., 5, 279 (1984)). Since these studies relate to proteins in organisms and the above-described method cannot inhibit the Maillard reaction that is produced in organisms, it has been hoped that a novel technique might be devleoped.

Meanwhile, GB2159408A (U.S.SN 712819) discloses a humectant which comprises (A) a silcion compound having at least one silanol group in the molecular thereof and (B) a water-soluble polymer substance which is most suitably a mucopolysaccharide, protein, cellulose derivative, starch derivative or polyvinyl alcohol.

Summary of the Invention

The inventors focused their attention on collagen of the skin and investigated the inhibition effect of collagen on the Maillard reaction (aged crosslinking) using various compounds. As a result, he achieved the present invention on the basis of the finding that a silicon compound having silanol groups in its molecule exhibits the above-described effect.

It is therefore a primary object of the present

invention to provide a novel technique which is capable of effectively inhibiting the Maillard reaction produced in organisms, particularly the crosslinking reaction of collagen. Another object of the present invention is to provide a technique which is capable of inhibiting the crosslinking reaction of collagen so as to inhibit the senescence of connective tissue.

These and other objects of the present invention will be clear from the following description.

In accordance with the present invention, there is provided a crosslinking reaction inhibitor for collagen, which comprises a silicon compound having at least one silanol group in its molecule.

Brief Description of the Drawings

Fig. 1 shows the gel electrophoresis of the CB peptides of collagen; and

Fig. 2 shows the densitometry of the gel electrophoresis.

Description of the Preferred Embodiments

The silicon compound used in the present invention may be any compound having at least one silanol group in its molecule, and it is preferable to use a compound having the formula (I):



wherein R is an alkyl, aralkyl, alkenyl, cycloalkyl, or alkoxy

0242855

gorup of 1 to 12 carbon atoms, preferably 1 to 6 carbon atoms,
two or more of R are the same or different from each other,
and n is an integer of between 0 and 3, preferably between 1
and 3 or a dehydrating condensation product thereof. Examples
5 of such silicon compounds include one or mixtures of two or
more of methyl silanetriol, dimethyl silanediol, trimethyl
silanol, ethyl silanetriol, diethyl silanediol, propyl
silanetriol, octyl silanetriol, dodecyl silanetriol, benzyl
silanetriol, phenylethyl silanetriol, propenyl silanetriol,
10 cyclohexyl silanetriol, methoxy silanetriol, and ethoxy
silanetriol. It is also possible to use a dimer or trimer
produced by the dehydrating condensation of methyl
silanetriol. Among the above-described compounds, it is
particularly preferable in the present invention to use methyl
15 silanetriol.

The above-described silicon compounds having silanol
groups in their molecules can be produced by the following
method:

20 (i) The corresponding chlorosilanes are hydrolyzed and
reacted with alkali hydroxide to obtain siliconates which show
pH values adjusted to between 4 and 5 by a cation exchange
resin or various acids, or

25 (ii) the corresponding alkoxysilanes are hydrolyzed.

These silicon compounds can be used as an inhibitor

for the crosslinking formation of collagen as they are or in the state of being dissolved or dispersed in a solvent such as water or ethanol. In addition, it is possible to add various additives such as an antiseptic or a perfume to the inhibitor
5 for the formation of crosslinking.

The silicon compound used in the present invention involves no problem with respect to safety. Shown below are data as to the safety of the silicon compounds.

LD50

(mg/kg)

10 Methyl silanetriol

1,000 or more

(Rat subcutaneous treatment)

The present invention is capable of effectively
15 preventing the corsslinking of collagen by sugars. Therefore, the use of the inhibitor of the present invention is able to effectively prevent various types of senescence such as skin senescence produced by the crosslinking of collagen present in the human body.

20 Thus, the crosslinking inhibitor of the present invention can be used as a skin preparation such as a cosmetic or ointment, or in a medical treatment such as a method involving injection, or by adding it to a health drink or a health food.

25 When the inhibitor is used in a skin preparation, it is preferable that the preparation contain 0.01 to 10 wt% (referred to as % hereinafter) of the inhibitor of the present

invention. When the inhibitor is used for medical treatment by injection or as a food, it is preferable to use the inhibitor in an amount of 0.1 to 10 mg and 0.01 to 1 mg relative to 1 kg of body weight, respectively.

5 The present invention is described in detail below with reference to unlimitative examples.

Example 1

10 Methyl silanetriol was produced by the following method and the inhibitor of the present invention containing the methyl silanetriol produced was added to collagen or a system comprising collagen and glucose in various amounts to form various samples, and the effects thereof in inhibiting the corsslinking of collagen were examined.

15 Method of producing methylsilanetriol

 The precipitates produced by hydrolysis of methyltrichlorosilane were recovered and dissolved in a solution of sodium hydroxide in an amount which was 3 times
20 that of the precipitates to prepare an aqueous solution of trisodium methylsiliconate.

 This aqueous solution was diluted with water and the pH value thereof was adjusted to between 4 and 5 by a cation exchange resin or various acids to form an aqueous solution of
25 methyl silanetriol.

Method of measuring the effect of crosslinking inhibition

100 mg of collagen obtained by grinding and deliming the bone of a spring chicken, 1 ml of 0.2M phosphate buffer solution (pH 7.4) of 0.2M glucose, and a given amount of the
5 silicon compound were reacted at 37°C for 4 weeks. After the reaction was completed, the collagen was separated by filtering the reaction solution and 10 mg of the collagen was cut with bromine cyanide (CNBr) in accordance with a conventional method (the formation of CB peptides) and then
10 subjected to acrylamide gel electrophoresis.

An example of the pattern obtained from the gel electrophoresis of the CB peptides is shown in Fig. 1 and the result of the measurement by densitometry is shown in Fig. 2. Table 1 shows the ratios of portion A (a low-molecular weight
15 region) to portion B (a high-molecular weight region: high-molecular substances of collagen having crosslinking formed by the Maillard reaction) in Fig. 1 and the rates of the collagen solubilized by the cutting with CNBr, together with the compositions of the samples used.

20 In Fig. 1, the collagen migrates from the upper portion B toward the lower portion A of the figure, the upper portion B being the crosslinking region and the lower portion A being the non-crosslinking region. The respective bands are identified as the peptides shown on the right side of the
25 figure. In Fig. 2, the right side is the high-molecular weight region and the left side is the low-molecular weight region. Reference numbers in the figure correspond to those

0242855

of the main peaks of the gel electrophoresis in Fig. 1, they correspond to the following:

- 1 α 1CB8
- 2 α 2CB4 + α 1CB7
- 3 α 1CB3. 7
- 4 α 1CB6. 7
- 5 α 1CB3. 5

Table-1

Sample No.	Composition of sample			Result	
	Collagen (mg)	Glucose (M)	Methyl-silanetriol (%)	Portion A/portion B	Soluble collagen (%)
1	100	-	-	0.11	100
2	100	0.2	-	0.53	54
3	100	0.2	0.003	0.55	50
4	100	0.2	0.01	0.40	81
5	100	0.2	0.03	0.35	86
6	100	0.2	0.1	0.40	76
7	100	0.2	0.3	0.54	54

It can be seen from the above results that the addition of 0.01 to 0.1% of methyl silanetriol inhibits the crosslinking and polymerization of collagen and thus silanol compounds have the effect of inhibiting the crosslinking.


5 Since the concentration of collagen and glucose in the living body is lower than that of the evaluation method of the present invention, it is thought that the corsslinking takes place over a very long time and the proper amount of a silanol to be compound administered is not limited to the amount of
10 0.01 to 0.1% for which the effect in the present invention was demonstrated.

Example 2

One silicon compound was selected from the group
15 consisting of dimethy silanediol, trimethyl silanol, dodecyl silanetriol, cyclohexyl silanetriol, propenyl silanetriol and a dimer of methy silanetriol, and the effects thereof in inhibiting the crosslinking of collagen were measured by the same method as set forth in Example 1. The results obtained
20 are shown in Table 2.

Table 2

Sample No.	Composition of sample			Result	
	Collagen (mg)	Glucose (M)	Silicon compound (%)	Portion A/ Portion B	Soluble collagen (%)
1	100	-	-	0.11	100
2	100	0.2	-	0.53	54
8	100	0.2	Dimethyl silanediol	0.43	76
9	100	0.2	0.1	0.35	80
10	100	0.2	Trimethyl silanol	0.45	70
11	100	0.2	0.1	0.38	75
12	100	0.2	Phenyl silanetriol	0.40	70
13	100	0.2	0.05	0.41	72
14	100	0.2	Dodecyl silanetriol	0.45	68
15	100	0.2	0.05	0.44	72
16	100	0.2	Cyclohexyl silanetriol	0.42	68
17	100	0.2	0.05	0.45	73
18	100	0.2	Propenyl silanetriol	0.45	71
19	100	0.2	0.1	0.39	72
20	100	0.2	Dimer of methyl silnetriol*)	0.35	80
21	100	0.2	0.05	0.40	75



Example 3

Various compositions to which the crosslinking inhibitor for collagen in the present invention was added are shown below.

Example of Cream Composition I

	Ingredient	Content wt part
Oil phase	Liquid paraffin	6
	Paraffin wax	1
	Isopropyl palmitate	3
	Spermaceti wax	2
	Cetyl alcohol	2
	Stearic acid	2
	Glycerin monostearate	1.45
	Polyoxyethylene (40) monostearate	1.35
	n-butyl-p-hydroxybenzoate	0.1
Water phase	Methyl p-hydroxybenzoate	0.2
	Aqueous solution of 1% methyl silanetriol	50
	Purified water	30.9

Example of Cream Composition II

	Ingredient	Content wt part
Oil phase	Liquid paraffin	6
	Squalane	3
	Cetyl palmitate	2
	Cetyl alcohol	2
	Stearic acid	2
	n-butyl-p-hydroxybenzoate	0.1
Water phase	Methyl p-hydroxybenzoate	0.1
	Aqueous solution of 10% dimethyl silamediol	50
	Purified water	34.8

Example of Cosmetic Lotion

Ingredient	Content wt part
Aqueous solution of 0.05% of phenyl silanetriol	40
Propylene glycol	5
Ethanol	5
Disodium hydrogenphosphate	0.05
Methyl p-hydroxybenzoate	0.1
Perfume	0.2
Purified water	49.65

Example of Ointment Composition

	Ingredient	Content wt part
Oil phase	White vaseline	25
	Stearyl alcohol	22
	n-butyl-p-hydroxybenzoate	0.015
Water phase	Propylene glycol	12
	Sodium dodecyl sulfate	1.5
	Methyl p-hydroxybenzoate	0.025
	Aqueous solution of 2% methyl silanetriol	25
	Purified water	14.46

Example of Ointment Composition

	Ingredient	Content wt part
Oil phase	White vaseline	40
	Cetyl alcohol	18
	Sorbitane sesquiolate	5
	Polyoxyethylene (9) dodecylalcohol	0.15
	n-butyl-p-hydroxybenzoate	0.1
Water phase	Methyl-p-hydroxybenzoate	0.1
	Aqueous solution of 0.05% cyclohexyl silanetriol	20
	Purified water	16.3

Example of Composition of Injection Solution

Ingredient	Content wt part
Aqueous solution of 1% methyl silanetriol	50
Salt	0.9
Purified water	49.1

Example of Composition of Health Drink

Ingredient	Content wt part
Aqueous solution of 1% methyl silanetriol	1
Sugar	10
Perfume	Suitable amount
Purified water	89

What is claimed is:

1. A crosslinking reaction inhibitor for collagen, which comprises a silicon compound having at least one silanol group in its molecule.

2. A crosslinking inhibitor according to Claim 1, wherein said silicon compound is a compound having the formula (I):



wherein R is an alkyl, aralkyl, alkenyl, cycloalkyl, or alkoxy group, two or more of R being the same or different from each other, and n is an integer of from 0 to 3 or a dehydrating condensation product thereof.

3. A crosslinking inhibitor according to Claim 1, wherein said silicon compound is methyl silanetriol.

4. A solution for injection containing a crosslinking reaction inhibitor for collagen as set forth in Claim 1.

5. Food containing a crosslinking reaction inhibitor for collagen as set forth in Claim 1.

6. A skin preparation containing a crosslinking reaction inhibitor for collagen as set forth in Claim 1.

FIG. 1





FIG. 2





Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 242 855
A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **87105854.1**

(51) Int. Cl.: **A 61 K 7/48, A 61 K 9/00,
A 23 L 1/03**

(22) Date of filing: **21.04.87**

(30) Priority: **21.04.86 JP 91769/86**

(71) Applicant: **LION CORPORATION, 3-7, Honjo 1-chome,
Sumida-ku Tokyo (JP)**

(43) Date of publication of application: **28.10.87
Bulletin 87/44**

(72) Inventor: **Nagaoka, Yoshio, No. 1-17-9-102,
Higashiyotsugi Katsushika-ku, Tokyo (JP)**
Inventor: **Yasumasu, Tomoko, No. 500-1-3-309,
Hongo-cho, Funabashi-shi Chiba-ken (JP)**
Inventor: **Noda, Seiji, No. 531-464, Wakamatsu-cho,
Chiba-shi Chiba-ken (JP)**
Inventor: **Higo, Moriaki, No. 6-8-8, Makuharinishi,
Chiba-shi Chiba-ken (JP)**


(84) Designated Contracting States: **AT BE CH DE ES FR GB
IT LI NL SE**

(88) Date of deferred publication of search
report: **07.09.88 Bulletin 88/36**

(74) Representative: **Kuhnen, Wacker & Partner,
Schneggstrasse 3-5 Postfach 1729, D-8050 Freising (DE)**

(54) **Crosslinking reaction inhibitor for collagen.**

(57) The present invention relates to a crosslinking inhibitor which inhibits the formation of crosslinking (aged crosslinking) of collagen by utilizing the Maillard reaction between collagen and sugar. The inhibitor comprises a silicon compound having at least one silanol group in its molecule, such as methyl silanetriol, dimethyl silanediol, trimethyl silanol, dodecyl silanetriol, cyclohexyl silanetriol or propenyl silanetriol.



EP 0 242 855 A3



European Patent
Office

EUROPEAN SEARCH REPORT

0242855

Application Number

EP 87 10 5854

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D, A	FR-A-2 561 915 (LION CORP.) * Page 3, lines 18-31; page 4, lines 25-29; page 6, lines 5-25; page 7, lines 33-34; examples 2,5,6; claims * -----	1-3,6	A 61 K 7/48 A 61 K 9/00 A 23 L 1/03
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K A 23 L 1/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 14-06-1988	Examiner ESPINOSA Y CARRETERO M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	